


Heating of coronal X-ray Bright Points

Biswajit Mondal

NASA Postdoctoral Program
NASA Marshall Space Flight Centre
Huntsville, Alabama, USA

Amy R. Wineberger (NASA Marshall Space Flight Centre),
P.S. Athiray (Univ. of Alabama in Huntsville, NASA MSFC)



Jan-17-2024

About me..



Biswajit Mondal

Graduated from Physical Research
Laboratory, India, 2022

NPP at NASA Marshall Space Flight
Center, USA, 2023

Contact me:
biswajit.mondal@nasa.gov

Group and collaborators



MSFC heliophysics team:

Dr. Amy R. Wineberger,

Dr. P.S. Athiray,

Dr. Sabrina L. Savage
and MaGIXS team.

Collaborators:

Dr. Helen E. Mason (Univ. Cambridge)



Dr. Giulio Del Zanna (Univ. Cambridge)

Dr. J.A. Klimchuk (NASA GSFC)

Dr. Aveek Sarker (PRL)

Dr. Santosh Vadawale (PRL) and others!

What are the XBPs?




- XBPs are located all over the solar disk. (Vaiana et al., 1973; Krieger et al., 1971; Golub et al., 1974)
- In the solar maxima, their contribution is hidden behind the huge AR emission.
- During the quiet phase XBPs are the primary on-disk X-ray contributors.

Questions

- Fraction of the total quiet Sun X-rays contributed by the XBPs
- Can nanoflare maintained the heating of the XBPs?
- If yes, what is the nanoflare frequency?


Differential Emission Measure (DEM) -> Diagnostics of Multi-thermal plasma



DEM reconstructions


$$O_i = \int_T DEM(T) R_i(T) dT + \delta O_i$$

Observed
Intensity






Chandrayaan-2 Solar X-ray Monitor (XSM)

- Disk-integrated soft X-ray spectrometer of the Sun
- Energy range: 1-15 keV(< M5 class) or 2-15 keV (> M5).
- Energy resolution: of 175 eV @ 5.9 keV.
- Cadence: 1 second
- Dynamic range: Sub-A to X-9 class flux level



XSM website: <https://www.prl.res.in/ch2xsm/>


XSM Observations during the minimum of solar cycle 24



- XSM spectra has the informations of temperature, EM as well as the abundances of Mg, Al, and Si.
- Vadawale, et al, (ApJL 912:L13, 2021 & 912:L12)


Extracting the XER emission from AIA

AIA 193 A




e

XER on AIA 193 A




f

XRT Be thin




DEM of Full sun & XERs



Full Sun \rightarrow XER + diffuse corona

XER \rightarrow XBP's + limb

Separate the different QS component




$$\mathcal{R} = \sum_i EMD(T_i) \Lambda(T_i)$$


DEM used	$\mathcal{R}(5.6 \leq \log T \leq 6.1)$ ($\text{erg cm}^{-2} \text{s}^{-1}$)	$\mathcal{R}(6.1 \leq \log T \leq 6.4)$ ($\text{erg cm}^{-2} \text{s}^{-1}$)
$DEM_{FullSun}$	0.78×10^5	0.09×10^5
DEM_{XER}	1.69×10^5	1.01×10^5
DEM_{XBPs}	1.08×10^5	0.87×10^5

- Lower temperature radiation loss from the diffuse corona is significant whereas at higher temperature it is negligible.
- Lower temperature XBP's emission is more than 63%, while at higher temperature it is more than 85%







Can nanoflares be responsible to heat the XBPs?



- Modeled the XBP's assuming nanoflare heating scenario.
- Estimate the simulated DEM, which is compared with the observed DEM.
- We used EBTEL code (Klimchuk et al. 2008; Cargill et al. 2012; Barnes et al. 2016).



Magnetic skeleton of XBPs



$$N_i = \frac{A_i}{2\pi r^2}$$

assume, $r = 1 \text{ Mm}$

Heating function

- Here we consider that nanoflares occur with the release of stored magnetic energy (Parker, 1988)
- Magnetic stored energy density :
$$E = \frac{(\tan(\theta) < B >)^2}{8\pi} (erg\ cm^{-3})$$
- $\tan(\theta) = c \rightarrow 0.2 - 0.3$, to satisfy observed coronal heating requirement (Parker 1988; Klimchuk 2015).
- Consider triangular heating profiles having a duration (τ) of 100 s.
- The peak heating rate during an event is randomly chosen between minimum (H_0^{min}) and maximum (H_0^{max}) values that are loop dependent.

$$H_{0_{ij}}^{max} = \frac{1}{\tau} \frac{(c < B >_{ij})^2}{8\pi} (erg\ cm^{-3}\ s^{-1})$$

$$H_0^{min} = 0.01 H_0^{max}$$

$$d_{ij}^l = \frac{\tau L}{F} \times H_{ij}^{l-1}$$


H_0 is randomly chosen between minimum (H_{0ij}^{min}) and maximum (H_{0ij}^{max}) for a loop.

- Model parameters $\rightarrow c$ and F


Heating function

Coronal radiation loss


- Constant F : $F = g \times \boxed{1.95 \times 10^5 \text{ (erg cm}^{-2} \text{ s}^{-1})}$; $g \rightarrow 2 \text{ to } 3$ (Klimchuk et al. 2008)
- Variable F : $F_{ij} = -\frac{1}{4\pi} V_h \tan(\theta) B_{ij}^{base} < B >_{ij}$; $V_h \rightarrow 0.5 \text{ to } 2 \text{ km/s}$



$L = 30 \text{ Mm}$
 $< B > = 10 \text{ G}$
 $c = 0.25$
 $g = 2.0$
 $V_h = 1 \text{ km/s}$
 $B_{base} = 15 \text{ G}$




Simulated DEM convolved with instrument responses



- Still, model DEM predicts 2-3 times higher emission at lower T.
- Simulated TR predicts a larger emission than the observed one → common (e.g., Warren et al. 2008) in loop simulations.
- Possibility – absorption TR emission by frequent chromospheric jets, such as spicules (De Pontieu et al. 2009)

➤ Agreement for the coronal portion of the loops suggest nanoflare can maintain the heating of XBP.


Frequency distribution with nanoflare energy



- Power-law slope -2.5 : Nanoflares energy > larger flares or microflares.

Detailed study for a single XBP

Marshall Grazing Incident X-ray Spectrometer (MaGIXS)






Champey et al. 2022




July-30-2021, 18:20 UT


Overlapping spectral images observed by MaGIXS




Hydrodynamic simulation of XBP-1




Used 1D
hydrodynamic
code, HYDRAD



Comparison between simulations and observations




Results & Discussions



- HF nanoflare -> average delay = 1500 - 3000 s (Which is smaller than the cooling time of the loops derived from the formula by Cargill et al 2014).
- Average Poynting flux = $\sim 5 \times 10^5$ erg/cm²/s
- MaGIXS and XRT are more sensitive.


Observed and Predicted images of XBP-1

Observed by AIA & XRT



Simulated

Derived from MaGIXS




Simulated

Summary

- Most of the high temperature emissions of the Sun in the absence of ARs comes from XBPs
- Nanoflare can maintained the heating of XBPs (average temperature of 2MK).
- Nanoflares delay time is in the range of 1500 s to 3000 s.
- The average poynting flux of XBPs are found to be around $5 \times 10^5 \text{ erg cm}^{-2} \text{ s}^{-1}$.
- MaGIXS and XRT are more sensitive to diagnose the heating frequency, whereas only AIA channels are less sensitive.

MaGIXS-2 (This year)



Thank You for your attention!